

Revised: February 2017 (6th version)

Standard Commodity Classification No. of Japan
875200

-Kampo-preparation-

Powerful drug **SANWA Hachimijoganryo Extract Fine Granules**

&lt;hachimijoganryo&gt;

S-04

This product is easy to take, fine granules made from the extract of "hachimijogan" listed in "KINKIYORYAKU".

**Storage. Precautions for handling**

Since this product is hygroscopic property, keep a container tightly closed in a cool place away from direct rays after use.

**Expiration date**

Use before the expiration date indicated on the label or the package.

Approval No.	(61AM)3622
Date of listing in the NHI reimbursement price	October 1986
Date of initial marketing in Japan	November 1986

**DESCRIPTION****Composition**

9g/day of this product contains 6.0g of a dried JP Hachimijogan extract of the following mixed crude drugs.

JP Rehmannia Root.....	5.0g
JP Cornus Fruit.....	3.0g
JP Dioscorea Rhizome.....	3.0g
JP Alisma Rhizome.....	3.0g
JP Poria Sclerotium.....	3.0g
JP Moutan Bark.....	3.0g
JP Cinnamon Bark.....	1.0g
JP Processed Aconite Root.....	1.0g

(JP: The Japanese Pharmacopoeia)

It also contains Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid as inactive ingredients.

**Product Description**

This product is fine granules, brown color, it has characteristic smell, and it tastes bitter and a little sweet.  
ID code: S-04

**INDICATIONS**

The following symptoms of those patients who have soft and weak abdomen, coldness and pain in the low back, decreased urinary output or increased urinary frequency, hot feeling in the body or the extremities

Chronic nephritis, diabetes mellitus, edema, edema by beriberi, bladder catarrh, low back pain, frozen shoulder and shoulder stiffness

**DOSAGE AND ADMINISTRATION**

The usual adult dose is 9g/day orally in 3 divided doses before or between meals. The dosage may be adjusted according to the patient's age and symptoms.

**PRECAUTIONS****1. Careful Administration (Hachimijoganryo should be administered with care in the following patients.)**

(1)Patients with strong constitution[Adverse reactions are likely to occur, and the symptoms may be aggravated.]

- (2) Patients with sensitivity to heat, a tendency towards hot flush and red face. [Palpitation, hot flush, numbness of the tongue, nausea, etc. may occur.]
- (3) Patients with an extremely weak gastrointestinal tract [Anorexia, epigastric distress, nausea, vomiting, abdominal pain, diarrhea, constipation, etc. may occur.]
- (4) Patients with anorexia, nausea or vomiting [These symptoms may be aggravated.]

**2. Important Precautions**

- (1) When this product is used, the patient's "SHO" (constitution /symptoms) should be taken into account. The patient's progress should be carefully monitored, and if no improvement in symptoms/findings is observed, continuous treatment should be avoided.
- (2) When this product is coadministered with other Kampo-preparations (Japanese traditional herbal medicines),etc., attention should be paid to the duplication of the contained crude drugs. Special caution should be exercised when this product is coadministered with preparations containing Aconite Root. SHO: The term "SHO" refers to a particular pathological status of a patient evaluated by the Kampo diagnosis, and is patterned according to the patient's constitution, symptoms, etc. Kampo-preparations (Japanese traditional herbal medicines) should be used after confirmation that it is suitable for the identified "SHO" of the patient.

**3. Adverse Reactions**

This product has not been investigated (drug use investigations, etc.) to determine the incidence of adverse reactions. Therefore, the incidence of adverse reactions is not known.

	Incidence unknown
<b>Hypersensitivity</b> <sup>Note1)</sup>	Rash, Redness, Pruritus, etc.
<b>Hepatic</b>	Abnormality of hepatic function [Increased AST (GOT), ALT(GPT),and total bilirubin levels, etc.]
<b>Gastrointestinal</b>	Anorexia, Epigastric distress, Nausea, Vomiting, Abdominal pain, Diarrhea, Constipation, etc.
<b>Others</b>	Palpitation, Hot flush, Numbness of the tongue, etc.

Note1) If such symptoms are observed, administration should be discontinued.

**4. Use in the Elderly**

Because elderly patients often have reduced physiological function, careful supervision and measures such as reducing the dose are recommended.

**5. Use during Pregnancy, Delivery or Lactation**

Use of this product in pregnant women, women who may possibly be pregnant is not recommended. [Moutan Bark contained in this product may cause premature birth or abortion. Besides, adverse reactions due to Processed Aconite Root contained in this product are likely to occur.]

**6. Pediatric Use**

This product should be administered with care in children.  
[This product contains Processed Aconite Root.]

**PACKAGING**

Bottles of 500g  
3.0g x 300 packets

**Manufactured and Distributed by:**

Sanwa Shoyaku Co., Ltd.  
6-1, Hiraide-Kogyo-Danchi, Utsunomiya-city, Tochigi, 321-0905,  
Japan