

Revised: August 2009 (2nd version)

Standard Commodity Classification No. of Japan
875900

-Crude drug-

The Japanese Pharmacopoeia **Powdered Processed Aconite Root**Powerful drug **Kakobushimatsu 「SANWA SHOYAKU」**

<kakobushimatsu>

S - 01

This product is easy to take, powder made from Japanese Aconite Root which processed with high-pressure steam.

Storage • Caution

Keep a container tightly closed in a dry area away from light and heat.

Expiration date

Use before the expiration date indicated on the label or the outer package.

Approval No.	22000AMX02002000
Date of listing in the NHI reimbursement price	December 2008
Date of initial marketing in Japan	December 2008

COMPOSITION

Kakobushimatsu is powder made from Japanese Aconite Root (*Aconitum carmichaeli* Debeaux., or *Aconitum japonicum* Thunb.) which processed with high-pressure steam. It contains JP Corn Starch as an inactive ingredient.

(JP: The Japanese Pharmacopoeia)

DESCRIPTION

Kakobushimatsu is powder, light brown color, it has characteristic smell.

INDICATIONS

Cardiotonic, analgesia, and diuresis

DOSAGE AND ADMINISTRATION

The usual adult dose is 0.5g~1.5g/day orally. Kakobushimatsu blends with other Kampo-preparations (Japanese traditional herbal medicines).

PRECAUTIONS**1. Careful Administration (Kakobushimatsu should be administered with care in the following patients.)**

- (1) Patients with strong constitution [Adverse reactions are likely to occur, and the symptoms may be aggravated.]
- (2) Patients with sensitivity to heat, a tendency towards hot flush and red face. [Palpitation, hot flush, numbness of the tongue, nausea, etc. may occur.]

2. Important Precautions

When this product is coadministered with preparations containing Aconite Root, attention should be paid to the duplication.

3. Adverse Reactions

This product has not been investigated (drug use investigations, etc.) to determine the incidence of adverse reactions. Therefore, the incidence of adverse reactions is not known.

	Incidence unknown
Others	Palpitation, Hot flush, Numbness of the tongue, Nausea, etc.

4. Use in the Elderly

Because elderly patients often have reduced physiological function, careful supervision and measures such as reducing the dose are recommended.

5. Use during Pregnancy, Delivery or Lactation

The safety of this product in pregnant women has not been established. Therefore, use of this product in pregnant women, women who may possibly be pregnant is not recommended.

6. Pediatric Use

The safety of this product in children has not been established. [Insufficient clinical data.]

PACKAGING

Bottles of 500g
0.5g × 300 packets
0.5g × 1200 packets

Manufactured and Distributed by:

SANWA SHOYAKU CO., LTD.

6-1, Hiraide-Kogyo-Danchi, Utsunomiya-city, Tochigi 321-0905, Japan