

Sanwa Shoyaku Co., Ltd.

Revised: September 2014 (3rd version)

Standard Commodity Classification No. of Japan
875900

-Crude drug-

The Japanese Pharmacopoeia **Powdered Processed Aconite Root**Powerful drug **Kakobushimatsu [SANWA SHOYAKU]**

&lt;kakobushimatsu&gt;

S-01

This product is easy to take, powder made from Japanese Aconite Root which processed by autoclaving.

**Storage. Precautions for handling**

Since this product is hygroscopic property, keep a container tightly closed in a cool place away from direct rays after use.

**Expiration date**

Use before the expiration date indicated on the label or the package.

Approval No.	22000AMX02002000
Date of listing in the NHI reimbursement price	December 2008
Date of initial marketing in Japan	December 2008 September 1981 <sup>Note)</sup>

Note) Date of initial marketing by former brand name.

**DESCRIPTION****Composition**

This product is powder made from Aconite Root (*Aconitum carmichaeli* Debeaux., or *Aconitum japonicum* Thunb.) which processed by autoclaving(Process 1).

It contains Corn Starch as an inactive ingredient.

**Product Description**

This product is light brown color and it has characteristic smell.  
ID code: S-01

**INDICATIONS**

Cardiotonic, analgesia, and diuresis

**DOSAGE AND ADMINISTRATION**

The usual adult dose is 0.5g-1.5g/day orally with other Kampo-preparations(Japanese traditional herbal medicines).

**PRECAUTIONS****1. Careful Administration (Kakobushimatsu should be administered with care in the following patients.)**

- (1)Patients with strong constitution[Adverse reactions are likely to occur, and the symptoms may be aggravated.]
- (2)Patients with sensitivity to heat, a tendency towards hot flush and red face. [Palpitation, hot flush, numbness of the tongue, nausea, etc. may occur.]

**2. Important Precautions**

When this product is coadministered with preparations containing Aconite Root, attention should be paid to the duplication.

**3. Adverse Reactions**

This product has not been investigated (drug use investigations, etc.) to determine the incidence of adverse reactions. Therefore, the incidence of adverse reactions is not known.

	<b>Incidence unknown</b>
<b>Others</b>	Palpitation, Hot flush, Numbness of the tongue, Nausea, etc.

**4. Use in the Elderly**

Because elderly patients often have reduced physiological function, careful supervision and measures such as reducing the dose are recommended.

**5. Use during Pregnancy, Delivery or Lactation**

The safety of this product in pregnant women has not been established. Therefore, use of this product in pregnant women, women who may possibly be pregnant is not recommended.

**6. Pediatric Use**

The safety of this product in children has not been established.  
[Insufficient clinical data.]

**PACKAGING**

Bottles of 500g  
1.5g x 300 packets  
1.5g x 1,200 packets

**Manufactured and Distributed by:**

Sanwa Shoyaku Co., Ltd.  
6-1, Hiraide-Kogyo-Danchi, Utsunomiya-city, Tochigi, 321-0905, Japan