

Revised: January 2017 (5th version)

Standard Commodity Classification No. of Japan
875200

-Kampo-preparation-

SANWA Hangeshashinto Extract Fine Granules<hangeshashinto>
S-18

This product is easy to take, fine granules made from the extract of "hangeshashinto" listed in "SHOKANRON".

Storage. Precautions for handling

Since this product is hygroscopic property, keep a container tightly closed in a cool place away from direct rays after use.

Expiration date

Use before the expiration date indicated on the label or the package.

Approval No.	(61AM)3598
Date of listing in the NHI reimbursement price	October 1986
Date of initial marketing in Japan	November 1986

CONTRAINDICATIONS (Hangeshashinto is contraindicated in the following patients.)

1. Patients with aldosteronism
 2. Patients with myopathy
 3. Patients with hypokalemia
- [1-3: These diseases or symptoms may be aggravated.]

DESCRIPTION**Composition**

7.5g/day of this product contains 4.9g of a dried JP Hangeshashinto extract of the following mixed crude drugs.

JP Pinellia Tuber.....	5.0g
JP Scutellaria Root.....	2.5g
JP Processed Ginger.....	2.5g
JP Ginseng.....	2.5g
JP Glycyrrhiza.....	2.5g
JP Jujube.....	2.5g
JP Coptis Rhizome.....	1.0g

(JP: The Japanese Pharmacopoeia)

It also contains Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid as inactive ingredients.

Product Description

This product is fine granules, yellow-brown color, it has characteristic smell, and it tastes bitter and a little sweet.

ID code: S-18

INDICATIONS

The following symptoms of those patients with blocked feeling in the stomach, nausea, vomiting, coated tongue, feeling of residual fluid in the stomach, anorexia, borborygmus, and diarrhea or loose stools:

Acute or chronic gastrointestinal catarrh, fermentative diarrhea, stomatitis, dyspepsia, gastroparesis, gastric atony, mild illness or prognosis of gastric and duodenal ulcer, and hyperemesis gravidarum

DOSAGE AND ADMINISTRATION

The usual adult dose is 7.5g/day orally in 3 divided doses before or between meals. The dosage may be adjusted according to the patient's age and symptoms.

PRECAUTIONS**1. Important Precautions**

- (1)When this product is used, the patient's "SHO" (constitution /symptoms) should be taken into account. The patient's progress should be carefully monitored, and if no improvement in symptoms/findings is observed, continuous treatment should be avoided.
- (2) Since this product contains Glycyrrhiza, careful attention should be paid to the serum potassium level, blood pressure, etc., and if any abnormality is observed, administration should be discontinued.
- (3) When this product is coadministered with other Kampo-preparations (Japanese traditional herbal medicines),etc., attention should be paid to the duplication of the contained crude drugs.

SHO: The term "SHO" refers to a particular pathological status of a patient evaluated by the Kampo diagnosis, and is patterned according to the patient's constitution, symptoms, etc. Kampo-preparations (Japanese traditional herbal medicines) should be used after confirmation that it is suitable for the identified "SHO" of the patient.

2. Drug Interactions

Precautions for coadministration (Hangeshashinto should be administered with care when coadministered with the following drugs.)

Drugs	Signs, Symptoms, and Treatment	Mechanism and Risk Factors
(1)Preparations containing Glycyrrhiza	Pseudoaldosteronism is likely to occur.	Since glycyrrhizinic acid and diuretics have an accelerating action on the potassium excretion at the renal tubules, an acceleration of decrease in the serum potassium level has been suggested.
(2)Preparations containing glycyrrhizinic acid or glycyrrhizates	Besides, myopathy is likely to occur as a result of hypokalemia. (Refer to the section "Clinically significant adverse reactions".)	
(3)Loop diuretics Furosemide Etacrynic acid		
(4)Thiazide diuretics Trichlormethiazide		

3. Adverse Reactions

This product has not been investigated (drug use investigations, etc.) to determine the incidence of adverse reactions.

Therefore, the incidence of adverse reactions is not known.

(1) Clinically significant adverse reactions

- 1) **Interstitial pneumonia:** If fever, cough, dyspnea, abnormal pulmonary sound (fine crackle), etc. are observed, administration of this product should be discontinued, and examinations such as X-ray should be performed immediately and appropriate measures such as administration of adrenocortical hormones taken. Besides, the patient should be advised to discontinue this product immediately and to make contact with the physician in the event of fever, cough, dyspnea, etc.
- 2) **Pseudoaldosteronism:** Pseudoaldosteronism such as hypokalemia, increased blood pressure, retention of sodium/body fluid, edema, increased body weight, etc. may occur. The patient should be carefully monitored (measurement of serum potassium level, etc.), and if any abnormality is observed, administration should be discontinued and appropriate measures such as administration of potassium preparations should be taken.
- 3) **Myopathy:** Myopathy may occur as a result of hypokalemia. The patient should be carefully monitored, and if any abnormality such as weakness, convulsion/paralysis of limbs, etc. are observed, administration should be discontinued and appropriate measures such as administration of potassium preparations should be taken.
- 4) **Hepatic dysfunction and jaundice:** Hepatic dysfunction and/or jaundice with elevation of AST(GOT), ALT(GPT), AL-P and γ -GTP or other symptoms may occur. The patient should be carefully monitored for abnormal findings. Administration should be discontinued and appropriate therapeutic measures should be taken, if abnormalities are observed.

(2) Other adverse reactions

	Incidence unknown
Hypersensitivity ^{Note1)}	Rash, Urticaria, etc.

Note1) If such symptoms are observed, administration should be discontinued.

4. Use in the Elderly

Because elderly patients often have reduced physiological function, careful supervision and measures such as reducing the dose are recommended.

5. Use during Pregnancy, Delivery or Lactation

The safety of this product in pregnant women has not been established. Therefore, the product should be used in pregnant women, women who may possibly be pregnant only if the expected therapeutic benefits outweigh the possible risks associated with treatment.

6. Pediatric Use

The safety of this product in children has not been established.
[Insufficient clinical data.]

7. Precautions concerning Use

When taking this drug for stomatitis, the drug can be slowly swallowed after keeping it within the mouth.

PACKAGING

Bottles of 500g
2.5g x 300 packets

Manufactured and Distributed by:

Sanwa Shoyaku Co., Ltd.
6-1, Hiraide-Kogyo-Danchi, Utsunomiya-city, Tochigi, 321-0905, Japan