

Revised: February 2014 (3rd version)

Standard Commodity Classification No. of Japan
875200

-Kampo-preparation-

**SANWA Goreisanryo Extract Fine Granules**<goreisanryo>  
S-33

This product is easy to take, fine granules made from the extract of "goreisan" listed in "SHOKANRON" "KINKIYORYAKU".

**Storage. Precautions for handling**

Since this product is hygroscopic property, keep a container tightly closed in a cool place away from direct rays after use.

**Expiration date**

Use before the expiration date indicated on the label or the package.

Approval No.	(61AM)3629
Date of listing in the NHI reimbursement price	October 1986
Date of initial marketing in Japan	November 1986

**DESCRIPTION****Composition**

7.5g/day of this product contains 3.8g of a dried Goreisanryo extract of the following mixed crude drugs.

JP Alisma Rhizome.....	6.0g
JP Polyporus Sclerotium.....	4.5g
JP Poria Sclerotium.....	4.5g
JP Atractylodes Rhizome.....	4.5g
JP Cinnamon Bark.....	3.0g

(JP: The Japanese Pharmacopoeia)

It also contains Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid as inactive ingredients.

**Product Description**

This product is fine granules, brown color, it has characteristic smell, and it tastes bitter.

ID code: S-33

**INDICATIONS**

The following symptoms of those patients with dry mouth, dizziness, headache and edema, etc.:

Acute gastrointestinal catarrh, nausea, and nephrosis

**DOSAGE AND ADMINISTRATION**

The usual adult dose is 7.5g/day orally in 3 divided doses before or between meals. The dosage may be adjusted according to the patient's age and symptoms.

**PRECAUTIONS****1. Important Precautions**

- (1) When this product is used, the patient's "SHO" (constitution /symptoms) should be taken into account. The patient's progress should be carefully monitored, and if no improvement in symptoms/findings is observed, continuous treatment should be avoided.
- (2) When this product is coadministered with other Kampo-preparations (Japanese traditional herbal medicines), etc., attention should be paid to the duplication of the contained crude drugs.  
SHO: The term "SHO" refers to a particular pathological status of a patient evaluated by the Kampo diagnosis, and is patterned

according to the patient's constitution, symptoms, etc. Kampo-preparations (Japanese traditional herbal medicines) should be used after confirmation that it is suitable for the identified "SHO" of the patient.

**2. Adverse Reactions**

This product has not been investigated (drug use investigations, etc.) to determine the incidence of adverse reactions. Therefore, the incidence of adverse reactions is not known.

	Incidence unknown
<b>Hypersensitivity</b> <sup>Note1)</sup>	Rash, Redness, Pruritus, etc.
<b>Hepatic</b>	Abnormality of hepatic function [Increased AST(GOT), ALT(GPT) and $\gamma$ -GTP etc. ]

Note1) If such symptoms are observed, administration should be discontinued.

**3. Use in the Elderly**

Because elderly patients often have reduced physiological function, careful supervision and measures such as reducing the dose are recommended.

**4. Use during Pregnancy, Delivery or Lactation**

The safety of this product in pregnant women has not been established. Therefore, the product should be used in pregnant women, women who may possibly be pregnant only if the expected therapeutic benefits outweigh the possible risks associated with treatment.

**5. Pediatric Use**

The safety of this product in children has not been established. [Insufficient clinical data.]

**PACKAGING**

Bottles of 500g  
2.5g x 300 packets

**Manufactured and Distributed by:**

Sanwa Shoyaku Co., Ltd.

6-1, Hiraide-Kogyo-Danchi, Utsunomiya-city, Tochigi, 321-0905, Japan