

Revised: December 2014 (8th version)

Standard Commodity Classification No. of Japan
875900

-Crude drug-

## Aconinsan Tablets

ST-01

This product is the tablet made from Powdered Processed Aconite Root.

### Storage. Precautions for handling

Since this product is hygroscopic property, keep a container tightly closed in a cool place away from direct rays after use.

### Expiration date

Use before the expiration date indicated on the label or the package.

Approval No.	(40A)3987
Date of listing in the NHI reimbursement price	June 1967
Date of initial marketing in Japan	July 1967

## DESCRIPTION

### Composition

Active ingredient(per tablet) <sup>Note1)</sup>	Inactive ingredients
JP Powdered Processed Aconite Root (process1 <sup>Note2)</sup> 166.67mg	JP Lactose Hydrate or JP Corn Starch

Note1) 9 tablets/day of Aconinsan correspond to 1.5g of Powdered Processed Aconite Root.

Note2) Process1:Autoclaving.  
(JP: The Japanese Pharmacopoeia)

### Product Description

This product is the tablet, light brown color, it has characteristic smell, and it tastes bitter.

Diameter	Thickness	Weight
8.0mm	4.0mm	200mg

ID code: ST-01

## INDICATIONS

Analgesia, cardiotonic, and diuresis

## DOSAGE AND ADMINISTRATION

The usual adult dose is 9 tablets/day orally in 3 divided doses.

## PRECAUTIONS

### 1. Careful Administration ( Aconinsan should be administered with care in the following patients.)

- (1)Patients with strong constitution[Adverse reactions are likely to occur, and the symptoms may be aggravated.]
- (2)Patients with sensitivity to heat, a tendency towards hot flush and red face. [Palpitation, hot flush, numbness of the tongue, nausea, etc. may occur.]

### 2. Important Precautions

When this product is coadministered with preparations containing Aconite Root, attention should be paid to the duplication.

### 3. Adverse Reactions

This product has not been investigated (drug use investigations, etc.) to determine the incidence of adverse reactions. Therefore, the incidence of adverse reactions is not known.

Incidence unknown	
Others	Palpitation, Hot flush, Numbness of the tongue, Nausea, etc.

### 4. Use in the Elderly

Because elderly patients often have reduced physiological function, careful supervision and measures such as reducing the dose are recommended.

### 5. Use during Pregnancy, Delivery or Lactation

The safety of this product in pregnant women has not been established. Therefore, use of this product in pregnant women, women who may possibly be pregnant is not recommended.

### 6. Pediatric Use

The safety of this product in children has not been established. [Insufficient clinical data.]

## PHYSICOCHEMISTRY

JP Powdered Processed Aconite Root is powder made from Aconite Root(*Aconitum carmichaeli* Debeaux., and *Aconitum japonicum* Thunb.) which processed by autoclaving and reduced the toxicity. It contains Mesaconitine, Benzoylmesaconine, Benzoylaconine, etc. as ingredients.

## PRECAUTIONS FOR HANDLING

Since this product is hygroscopic property, keep a container tightly closed in a cool place away from direct rays after use.

## PACKAGING

900 tablets(3 tablets x 300 packets)  
Bottles of 1,000 tablets  
Bottles of 3,000 tablets

## Manufactured and Distributed by:

Sanwa Shoyaku Co., Ltd.  
6-1, Hiraide-Kogyo-Danchi, Utsunomiya-city, Tochigi, 321-0905, Japan